	FORM				PART	MEN	T OF CC	MMERC	E PATENT	AND TR.	ADEMARK	ATTO	RNEY'S DOCKET NUMBER X-11650
	1	•			L I.	ET	TER '	TO TI	HE UNI	TED S	STATES	U.S. A	APPLICATION NO. (if known, see 37 C.F.R. 1.5)
	_	D	ESI	GNAT	ED/	EL	ECTE	D OF	FICE (DO/EC	D/US)		10/01000
									DER 3	5 U.S.C	C. 371	IC DATE	10/018099
	IN	TER.		IONAL CT/US				NO.			NAL FILIN 000 (08.0		PRIORITY DATE CLAIMED 15 July 1999 (15.07.99)
	TITL	E O		VENTIO				ESS FO					DEPSIPEPTIDES
,	APPI	LICA	NT((S) FOR	DO/I	O/L	S: Ad	am Jose	eph Kreu	zman, P	Palaniappa ted Office (n Kulanth	haivel and Michael John Rodriguez (S) the following items and other
	infor			ewith Sub	mis	to m	e Onic	u State	s Designa	iicu/Licc	ica Office (DOIDOIO	by the following foliae and other
÷	1/[x	This	is a FIR	ST s	ubm	ission (of items	concerni	ng a filin	ig unde <u>r 35</u>	U.S.C. 37	<u>71</u>
	2.	\neg	/										ling under 35 U.S.C. 371.
	3.		This	express	requ	est to	begin	nationa	ıl examina	ation pro	cedures (35	5 U.S.C. 3	71(f)) at any time rather than delay
			exai	nination	until	the e	expirati	on of th	e applica	ble time	limit set in	35 U.S.C.	371(b) and PCT Articles 22 and 39(1).
	4.	X	A p	roper De	nand	for	Interna	tional P	reliminar	y Exami	nation was	made by tl	he 19th month from the earliest claimed
			prio	rity date.									
	5.	X	A c	opy of th	e Inte	rnat	ional A	pplicati	ion as file	d (35 U.	S.C. 371(c))(2))	
			a.	is tr	ansm	itted	l herew	ith (req	uired only	y if not tr	ansmitted l	by the Inte	rnational Bureau).
			b.						Internation				
			c.			_							ceiving Office (RO/US).
	6.										lish (35 U.S		
	7.	X	Am										cle 19 (35 U.S.C. 371 (c)(3))
			a.	are	trans	mitte	ed here	with (re	quired on	ly if not	transmitted	d by the Int	ternational Bureau).
			b.					•	e Internat				
			c.								for making	such amer	ndments has NOT expired.
			d.						ll not be r			10 (05 1)	G G 251()(2))
	8.		l									19 (35 U.:	S.C. 371(c)(3)).
	9.	X							or(s) (35 U				
	10.	X											ling any annexes, and, if not in English, an
		<u> </u>	Eng	glish lang	uage	tran	slation	of the a	nnexes to	the IPE	R under PC	I Article	36 (35 U.S.C. 371(c)(5)).
		ns 11	. to	16. belov	con	ceri	ı docui	ment(s)	or infor	mation i	ncluded:		
	11.	<u></u>	1								.97 and 1.9		
	12.	X	An	assignme	ent de	ocun	nent for	recordi	ing. A sep	parate co	ver sheet in	omplian	ce with 37 CFR 3.28 and 3.31 is included.
	13.	<u> </u>	ΑI	FIRST pr	elimi	nary	amend	ment.					
	1,4	<u></u>	AS	SECOND	or S	UBS	EQUE	NT pre	liminary a	amendme	ent.		
	14.	<u></u>	A s	substitute	spec	ifica	tion.						
	15.		A	hange of	pow	er of	attorn	ey and/o	or address	letter.			
	16.		Otl	ner items	or in	form	ation:				<u> </u>		

U.S. APPLICATION NO. (if	known, see 37 C.F.R. 1.5)	INTERNATIONAL PCT/U	APPLICATION NO. S00/15018	•	ATTORNEY'S DOC X-1	KET NUMBER 1650
	g fees are submitted:			CA	LCULATIONS	PTO USE ONLY
Neither internation nor international s	search fee (37 CFR 1.44	92(a)(1)-(5)): ation fee (37 CFR 1.482 45(a)(2)) paid to USPTC ared by the EPO or JPO)			
International preli	minary examination fee Search Report prepared	e (37 CFR 1.482) not pa by the EPO or JPO	id to USPTO \$890.00			
International preli	minary examination fee ch fee (37 CFR 1.445(a	e (37 CFR 1.482) not pa (2)) paid to USPTO	id to USPTO but \$740.00			
International prelibut all claims did	minary examination fee not satisfy provisions o	e paid to USPTO (37 Cl f PCT Article 33(1)-(4)	FR 1.482) \$710.00			
International preli 1.482) and all cla	iminary examination fee ims satisfied provisions	e paid to USPTO (37 Cl of PCT Article 33(2)-(RF 4) \$100.00			
	ENTER APPROP	RIATE BASIC FI	EE AMOUNT =	\$	890.00	
Surcharge of \$130.00 f	or furnishing the oath o	r declaration later than		\$		
months from the earlie	st claimed priority date NUMBER FILED	(37 CFR 1.492(e)). NUMBER EXTRA	RATE			L
Total claims	14 -20=	0	X \$18.00	\$		
Independent claims	7 - 3=	4	X \$84.00	\$	336.00	
	ENT CLAIM(S) (if app	olicable)	+ \$280.00	\$	280.00	
		OF ABOVE CAL	CULATIONS =	\$	616.00	
Reduction by 1/2 for fi		applicable. Verified Sm		\$	010.00	
	e 37 CFR 1.9, 1.27, 1.2			Ľ		
	000000000000000000000000000000000000000	1 1 1 1 1 1 1 1 1	SUBTOTAL =	\$	1,506.00	
Processing fee of \$130 months from the earlie	.00 for furnishing Englist claimed priority date	ish translation later than (37 CFR 1.492(f)).	12030+	\$		
-		TOTAL NAT	ΓΙΟΝΑL FEE =	\$	1,506.00	
Fee for recording the e	nclosed assignment (37 propriate cover sheet (3	CFR 1.21(h)). The ass	signment must be	\$		
accompanied by an app	propriate cover sheet (5	9	640.00 per property			
		TOTAL FEES	S ENCLOSED =	\$	1,506.00	
					Amount to be refunded	· ·
					charged	\$
b. X Please charg copy of this c. X The Commis	sheet is enclosed. ssioner is hereby author	to cover the above for No. 05-0840 in the amore tized to charge any additional and the days of this course of the course of	unt of <u>\$ 1,506.00</u> to tional fees which may			
to Deposit A	kecount ino. U5-U84U. <i>P</i>	A duplicate copy of this	SHOOL IS CHOIUSCU.			
NOTE: Where an ap (37 CFR 1.13	ppropriate time limit u 37(a) or (b)) must be f	nnder 37 CFR 1.494 or iled and granted to res	1.495 has not been store the application	met to p	, a petition to rev pending status. ー	ive
SEND ALL CORRESPO			Du gr	٠ ـ ١	tuel	
ELI LILLY AND COM PATENT DIVISION/11			SIGNAT	UKE	s	
LILLY CORPORATE O	CENTER		Tina M. ' NAM		ker	
INDIANAPOLIS, INDL	•	25885		نعه		
10 Dec 2	<u>u</u> PA	TENE TRADEMARK OFFICE	47,145 EGISTRATION NUMI	BER	(317) 277-3 TELEPHO	537 NE NUMBER
Bate		IPAGE 2 C				

20

diaminobutyric acid.

Rec'd PCT/PTO 11 DEC 2001

PROCESS FOR DEACYLATION OF LIPODEPSIPEPTIDES

FIELD OF THE INVENTION

The present invention relates to lipodepsipeptides, in particular, deacylation of the N-acyl side-chain of pseudomycin and syringomycin natural products and the compounds produced therefrom.

BACKGROUND OF THE INVENTION

Pseudomycins and syringomycins are natural products isolated from liquid cultures of *Pseudomonas syringae* (plant-associated bacterium) and have been shown to have antifungal activities. (see i.e., Harrison, L., et al., "Pseudomycins, a family of novel peptides from *Pseudomonas syringae* possessing broad-spectrum antifungal activity," <u>J. Gen. Microbiology</u>, **137**(12), 2857-65 (1991) and US Patent Nos. 5,576,298 and 5,837,685) Unlike the previously described antimycotics from *P. syringae* (e.g., syringomycins, syringotoxins and syringostatins), pseudomycins A-C contain hydroxyaspartic acid, aspartic acid, serine, dehydroaminobutyric acid, lysine and

The peptide moiety for pseudomycins A, A', B, B', C, C'

25 corresponds to L-Ser-D-Dab-L-Asp-L-Lys-L-Dab-L-aThr-Z-Dhb-L
Asp(3-OH)-L-Thr(4-Cl) with the terminal carboxyl group

closing a macrocyclic ring on the OH group of the N-terminal Ser. The analogs are distinguished by the N-acyl side chain, i.e., pseudomycin A is N-acylated by 3,4-dihydroxytetradeconoyl, pseudomycin A' by 3,4-dihydroxytetradecanoyl, pseudomycin B by 3-hydroxytetradecanoyl, pseudomycin B' by 3-hydroxydodecanoyl, pseudomycin C by 3,4-dihydroxyhexadecanoyl and pseudomycin C' by 3-hydroxyhexadecanoyl and pseudomycin C' by 3-hydroxyhexadecanoyl. (see i.e., Ballio, A., et al., "Novel bioactive lipodepsipeptides from Pseudomonas syringae: the pseudomycins," FEBS Letters, 355(1), 96-100, (1994) and Coiro, V.M., et al., "Solution conformation of the Pseudomonas syringae MSU 16H phytotoxic lipodepsipeptide Pseudomycin A determined by computer simulations using distance geometry and molecular dynamics from NMR data,"

Pseudomycins and syringomycins are known to have certain adverse biological effects. For example, destruction of the endothelium of the vein, destruction of tissue, inflammation, and local toxicity to host tissues have been observed when pseudomycin is administered intraveneously. Therefore, there is a need to identify compounds within this class that are useful for treating fungal infections without the currently observed adverse side effects.

Eur. J. Biochem., **257**(2), 449-456 (1998).)

5

WO 01/05815 PCT/US00/15018

BRIEF SUMMARY OF THE INVENTION

The present invention provides a process for deacylating the N-acyl side-chain of a lipodepsipeptide natural product to produce the corresponding nucleus. The deacylation of pseudomycin compounds produces the pseudomycin amino nucleus represented by the following structure I.

10 I

The nucleus is useful as a starting material for producing semi-synthetic derivatives of the corresponding natural product.

The process includes reacting a pseudomycin natural product with a deacylase enzyme selected from the group consisting of ECB deacylase and polymyxin acylase to produce

10

WO 01/05815 PCT/US00/15018

the corresponding nucleus represented by structure I. The free amine may rearrange to produce a cyclic peptide nucleus having a free hydroxy group represented by structure II below (also referred to as pseudomycin hydroxy nucleus).

$$H_2N$$
 H_2N
 H_2N
 H_2N
 H_2N
 H_2N
 H_2N
 H_2N
 H_3N
 H_4N
 H_4N
 H_5N
 H_5N
 H_5N
 H_7N
 H_7N

II

Compound II may then serve as starting material to generate novel derivatives which may be pharmaceutically active.

In another embodiment of the present invention, the process described above is used to deacylate syringomycin compounds to provide a syringomycin amino nucleus. For example, the amino nucleus of Syringomycin E has the following structure III.

III

Like the pseudomycin amino nucleus, the syringomycin amino nucleus may rearrange to form the following Compound IV (also referred to as syringomycin hydroxy nucleus).

IV

Even though specific chiral forms are depicted above for Compounds I, II, III and IV, other chiral forms are within the spirit of the present invention. Each of the compounds may also exist as pharmaceutically acceptable salts, hydrates or solvates thereof.

Definitions

As used herein, the term "pseudomycin" refers to compounds having the following formula:

where R is a lipophilic moiety. The lipophilic moiety includes C_9-C_{15} alkyl, C_9-C_{15} hydroxyalkyl, C_9-C_{15} dihydroxyalkyl, C_9-C_{15} alkenyl, C_9-C_{15} hydroxyalkenyl, or C_9-C_{15} dihydroxyalkenyl. The pseudomycin compounds A, A', B,

25

5

B', C, C' are represented by the formula I above where R is as defined below.

Pseudomycin A R = 3,4-dihydroxytetradecanoyl

Pseudomycin A' R = 3,4-dihydroxypentadecanoyl

Pseudomycin B R = 3-hydroxytetradecanoyl

Pseudomycin B' R = 3-hydroxydodecanoyl

Pseudomycin C R = 3,4-dihydroxyhexadecanoyl

Pseudomycin C' R = 3-hydroxyhexadecanoyl

DETAILED DESCRIPTION OF THE INVENTION

Applicants have discovered a process for enzymatically deacylating the N-acyl side-chain of a broad spectrum of lipodepsipeptide natural products to produce the corresponding nucleus. Surprisingly, the free amine nucleus rearranges to produce the free hydroxy derivative such as the compounds shown above as structures II and IV. Compounds I and III can be converted to Compounds II and IV, respectively, by exposing Compound I or III to a $pH \geq 6$. If the desired product is Compound I or III, then one could reduce the rate at which the rearranged product forms from the deacylated pseudomycin or deacylated syringomycin with the addition of an acid, such as trifluoroacetic acid. However, the addition of an acid could result in lower yields of the amine nucleus. At lower pHs, the enzyme may precipitate out of the reaction mixture thus stopping the

20

25

5

conversion. Therefore, the pH of the reaction mixture is preferably not lowered less than about 5.5. One could prevent enzyme precipitation by separating the enzyme from the reaction through a molecular weight membrane (i.e.,

10,000 to 50,000 molecular weight cutoff). The effluent through the membrane would contain compounds having a molecular weight less than 10,000 to 5,000 (e.g., Compounds I-IV) and would exclude the higher molecular weight enzyme. The effluent could then be pH adjusted down to stabilize the product.

Unlike acid deacylation processes (e.g., trifluoroacetic acid in an aqueous solvent at room temperature), the inventive enzymatic process may be used to deacylate pseudomycin or syringomycin analogs with or without gamma or delta hydroxy side chains. Therefore, the spectrum of starting natural products is expanded significantly. For example, one may deacylate pseudomycin A, A', B, B', C or C' using the inventive process. Whereas, the acid deacylation process is useful only with pseudomycin A, A' and C.

Suitable enzymes include ECB deacylase and Polymyxin acylase (available in both a crude & pure form as 161-16081 Fatty Acylase, Pure and 164-16081 Fatty Acylase, Crude, from Wako Pure Chemical Industries, Ltd.) ECB deacylase can be obtained from Actinoplanes utahensis (see e.g., LaVerne, D,

15

20

5

et al, "Deacylation of Echinocandin B by Actinoplanes utahensis," J. of Antibiotics, 42(3), 382-388 (1989).) The Actinoplanes utahensis ECB deacylase enzyme may be purified by the process described in U.S. Patent No. 5,573,936,

PCT/US00/15018

incorporated herein by reference. One may also use an enzyme that has been cloned and expressed in *Streptomyces lividans*. Attempts to deacylate pseudomycin A with Pen G Amidase and Phthalyl Amidase were not successful.

The enzymatic deacylation may be accomplished using standard deacylation procedures well known to those skilled in the art. For example, general procedures for using Polymyxin acylase may be found in Yasuda, N., et al, <u>Agric. Biol. Chem.</u>, 53, 3245 (1989) and Kimura, Y., et al., <u>Agric. Biol. Chem.</u>, 53, 497 (1989).

The deacylation process is generally ran at temperatures between about 20°C and about 60°C, preferably between about room temperature (25°C) and about 40°C. Higher temperatures may promote the formation of the rearranged product (Compound II). The enzyme is optimally active at pH 8.0 and at a temperature between about 50°C and 60°C. Although the reaction is faster at the higher pH and higher temperature, more rearranged product may be observed at the higher pH. Therefore, the pH of the reaction is

10

15

20

25

5

generally kept between about 5.5 and about 8.0. The reaction time will vary depending upon the pH and the temperature. However, with limiting enzyme concentration and saturated substrate concentration at high temperatures and pH, the reaction is linear through 10 minutes. Since Pseudomcyin A is unstable at higher pHs, deacylation of Pseudomycin A is generally ran at a lower pH (between about 5.0 and 6.0) and temperature (about 25°C). For example, deacylation of Pseudomycin A can be ran in a buffered solution containing 0.05 M KPO₄ and 0.8 M KCl. A saturated level of substrate is generally between about 0.5 mg and about 1 mg per ml of reaction.

As discussed earlier, pseudomycins are natural products isolated from the bacterium *Pseudomonas syringae* that have been characterized as lipodepsinonapetpides containing a cyclic peptide portion closed by a lactone bond and including the unusual amino acids 4-chlorothreonine (ClThr), 3-hydroxyaspartic acid (HOAsp), 2,3-dehydro-2-aminobutyric acid (Dhb), and 2,4-diaminobutyric acid (Dab). Methods for growth of various strains of *P. syringae* to produce the different pseudomycin analogs (A, A', B, B', C, and C') are generally described below and also described in more detail in PCT Patent Application Serial No. PCT/USOO/08728 filed by Hilton, et al. on April 14, 2000 entitled "Pseudomycin Production by *Pseudomonas Syringae*," incorporated herein by

20

5

reference, PCT Patent Application Serial No. PCT/US00/08727 filed by Kulanthaivel, et al. on April 14, 2000 entitled "Pseudomycin Natural Products," incorporated herein by reference, and U.S. Patent Nos. 5,576,298 and 5,837,685, each of which are incorporated herein by reference.

Isolated strains of P. syringae that produce one or more pseudomycins are known in the art. Wild type strain MSU 174 and a mutant of this strain generated by transposon mutagenesis, MSU 16H are described in U.S. Patent Nos. 5,576,298 and 5,837,685; Harrison, et al., "Pseudomycins, a family of novel peptides from Pseudomonas syringae possessing broad-spectrum antifungal activity, " J. Gen. Microbiology, 137, 2857-2865 (1991); and Lamb et al., "Transposon mutagenesis and tagging of fluorescent pseudomonas: Antimycotic production is necessary for control of Dutch elm disease," Proc. Natl. Acad. Sci. USA, 84, 6447-6451 (1987).

A strain of P. syringae that is suitable for production of one or more pseudomycins can be isolated from environmental sources including plants (e.g., barley plants, citrus plants, and lilac plants) as well as, sources such as soil, water, air, and dust. A preferred stain is isolated from plants. Strains of P. syringae that are isolated from environmental sources can be referred to as wild type.

10

20

25

5

used herein, "wild type" refers to a dominant genotype which naturally occurs in the normal population of P. syringae (e.g., strains or isolates of P. syringae that are found in nature and not produced by laboratory manipulation). Like most organisms, the characteristics of the pseudomycinproducing cultures employed (P. syringae strains such as MSU 174, MSU 16H, MSU 206, 25-B1, 7H9-1) are subject to variation. Hence, progeny of these strains (e.g., recombinants, mutants and variants) may be obtained by methods known in the art.

Mutant strains of P. syringae are also suitable for production of one or more pseudomycins. As used herein, "mutant" refers to a sudden heritable change in the phenotype of a strain, which can be spontaneous or induced by known mutagenic agents, such as radiation (e.g., ultraviolet radiation or x-rays), chemical mutagens (e.g., ethyl methanesulfonate (EMS), diepoxyoctane, N-methyl-Nnitro-N'-nitrosoguanine (NTG), and nitrous acid), sitespecific mutagenesis, and transposon mediated mutagenesis.

Pseudomycin-producing mutants of P. syringae can be produced by treating the bacteria with an amount of a mutagenic agent effective to produce mutants that overproduce one or more pseudomycins, that produce one pseudomycin (e.g., pseudomycin B) in excess over other pseudomycins, or that produce one or more pseudomycins under advantageous growth

5

conditions. While the type and amount of mutagenic agent to be used can vary, a preferred method is to serially dilute NTG to levels ranging from 1 to 100 μ g/ml. Preferred mutants are those that overproduce pseudomycin B and grow in minimal defined media.

Environmental isolates, mutant strains, and other desirable strains of P. syringae can be subjected to selection for desirable traits of growth habit, growth medium nutrient source, carbon source, growth conditions, amino acid requirements, and the like. Preferably, a pseudomycin producing strain of P. syringae is selected for growth on minimal defined medium such as N21 medium and/or for production of one or more pseudomycins at levels greater than about 10 μ g/ml. Preferred strains exhibit the characteristic of producing one or more pseudomycins when grown on a medium including three or fewer amino acids and optionally, either a lipid, a potato product or combination thereof.

Recombinant strains can be developed by transforming the *P. syringae* strains, using procedures known in the art. Through the use of recombinant DNA technology, the *P. syringae* strains can be transformed to express a variety of gene products in addition to the antibiotics these strains produce. For example, one can modify the strains to

WO 01/05815 PCT/US00/15018

introduce multiple copies of the endogenous pseudomycinbiosynthesis genes to achieve greater pseudomycin yield.

5

15

20

25

10

To produce one or more pseudomycins from a wild type or mutant strain of P. syringae, the organism is cultured with agitation in an aqueous nutrient medium including an effective amount of three or fewer amino acids, preferably glutamic acid, glycine, histidine, or a combination thereof. Alternatively, glycine is combined with one or more of a potato product and a lipid. Culturing is conducted under conditions effective for growth of P. syringae and production of the desired pseudomycin or pseudomycins. Effective conditions include temperatures from about 22°C to about 27°C, and a duration of about 36 hours to about 96 hours. Controlling the concentration of oxygen in the medium during culturing of P. syringae is advantageous for production of a pseudomycin. Preferably, oxygen levels are maintained at about 5 to 50% saturation, more preferably about 30% saturation. Sparging with air, pure oxygen, or gas mixtures including oxygen can regulate the concentration of oxygen in the medium.

Controlling the pH of the medium during culturing of P. syringae is also advantageous. Pseudomycins are labile at basic pH, and significant degradation can occur if the pH of the culture medium is above about 6 for more than about 12 hours. Preferably, the pH of the culture medium is

25

5

maintained between 6 and 4. *P. syringae* can produce one or more pseudomycins when grown in batch culture. However, fed-bath or semi-continuous feed of glucose and optionally, an acid or base (e.g., ammonium hydroxide) to control pH, enhances production. Pseudomycin production can be further enhanced by using continuous culture methods in which glucose and ammonium hydroxide are fed automatically.

Choice of *P. syringae* strain can affect the amount and distribution of pseudomycin or pseudomycins produced. For example, strains MSU 16H and 67 H1 each produce predominantly pseudomycin A, but also produce pseudomycin B and C, typically in ratios of 4:2:1. Strain 67 H1 typically produces levels of pseudomycins about three to five fold larger than are produced by strain MSU 16H. Compared to strains MSU 16H and 67 H1, strain 25-B1 produces more pseudomycin B and less pseudomycin C. Strain 7H9-1 are distinctive in producing predominantly pseudomycin B and larger amount of pseudomycin B than other strains. For example, this strain can produce pseudomycin B in at least a ten fold excess over either pseudomycin A or C.

As discussed earlier, the process described herein is also useful for deacylating syringomycin compounds.

Syringomycin E, syringotoxin B, and syringostatin A may be produced from cultures of *Pseudomonas syringae* pv. *syringae* strains B301D, PS268, and SY12, respectively. Syringomycin

5

A₁ and G may be isolated from *Pseudomonas syringae* pv. syringae as well. Strains B301D and PS268 are grown in potato dextrose broth as described by Zhang, L., and J. Y. Takemoto, "Effects of Pseudomonas syringae phytotoxin,

- Takemoto, "Effects of Pseudomonas syringae phytotoxin, syringomycin, on plasma membrane functions of Rhodotorula pilimanae," Phytopathol. 77(2):297-303 (1987). Strain SY12 was grown in syringomycin minimal medium supplemented with 100M arbutin (Sigma Chemical Co., A 4256; St. Louis, Mo.) and 0.1% fructose (SRMAF) (19, 23). SR-E, ST-B, and SS-A are purified by high performance liquid chromatography as described previously by Bidwai, A. P., and J. Y. Takemoto, "Bacterial phytotoxin, syringomycin, induces a protein kinase-mediatedphosphorylation of red beet plasma membrane polypeptides," Proc. Natl. Acad. Sci. USA, 84:6755-6759 (1987). Solubilized AmB containing 35% sodium deoxycholate (Sigma Chemical Co., A 9528; St. Louis, Mo.) and ketoconazole (Sigma Chemical Co., K-1003; St. Louis, Mo.)
- 20 lipodepsinonapeptides syringomycin E, syringotoxin B, and syringostatin A may be found in U.S. Patent No. 5,830,855, incorporated herein by reference.

production and isolation of three cyclic

The pseudomycin or syringomycin nucleus or corresponding rearranged compounds (Compounds II and IV) may be isolated and used per se or in the form of its

are used as test standards. A detailed description for the

5

pharmaceutically acceptable salt or solvate. The term "pharmaceutically acceptable salt" refers to non-toxic acid addition salts derived from inorganic and organic acids. Suitable salt derivatives include halides, thiocyanates, sulfates, bisulfates, sulfites, bisulfites, arylsulfonates, alkylsulfates, phosphonates, monohydrogen-phosphates, dihydrogenphosphates, metaphosphates, pyrophosphonates, alkanoates, cycloalkylalkanoates, arylalkonates, adipates, alginates, aspartates, benzoates, fumarates, glucoheptanoates, glycerophosphates, lactates, maleates, nicotinates, oxalates, palmitates, pectinates, picrates, pivalates, succinates, tartarates, citrates, camphorates, camphorsulfonates, digluconates, trifluoroacetates, and the like.

The term "solvate" refers to an aggregate that comprises one or more molecules of the solute (i.e., pseudomycin and syringomycin compound) with one or more molecules of a pharmaceutical solvent, such as water, ethanol, and the like. When the solvent is water, then the aggregate is referred to as a hydrate. Solvates are generally formed by dissolving the nucleus or rearranged compound (Compounds II or IV) in the appropriate solvent with heat and slowing cooling to generate an amorphous or crystalline solvate form.

20

WO 01/05815 PCT/US00/15018

EXAMPLES

Biological Samples

P. syringae MSU 16H is publicly available from the American Type Culture Collection, Parklawn Drive, Rockville, MD, USA as Accession No. ATCC 67028. P. syringae strains 25-B1, 7H9-1, and 67 H1 were deposited with the American Type Culture Collection on March 23, 2000 and were assigned the following Accession Nos.:

25-B1	Accession	No.	PTA-1622
7H9-1	Accession	No.	PTA-1623
67 H1	Accession	No.	PTA-1621

Chemical Abbreviations

The following abbreviations are used through out the examples to represent the respective listed materials:

ACN - acetonitrile

5

10

Ring of

15

25

TFA - trifluoroacetic acid

DMF - dimethylformamide

20 Example 1

Example illustrates the deacylation of Pseudomycin A using ECB Deacylase enzyme.

Pseudomycin A (50 μ g) and purified ECB Deacylase (50 μ l) in 900 μ l of an aqueous buffer solution containing 0.05 M potassium phosphate and 0.8 M potassium chloride. The pH remained between 6.0 and 8.0. The temperature was increased

10

15

20

from 25°C to 40°C. The reaction was monitored by HPLC (Waters C18 µBondapak 3.9 X 300 mm column, 235 nm, 1% acetonitrile/0.2% trifluoroacetic acid (4 minutes) to 60% acetonitrile/0.2% trifluoroacetic acid (16 minutes)). Both the pseudomycin amine nucleus (Compound I) and the rearranged pseudomycin hydroxy nucleus (Compound II) were observed.

Both Compounds ${\bf I}$ and ${\bf II}$ showed identical M+H ion (m/z 981.3) in the electrospray ionization mass spectroscopy (ESIMS) corresponding to a molecular formula of C₃₇H₆₁ClN₁₂O₁₇. (See Table I below) Detailed analysis of ¹H and TOCSY (total correlation spectroscopy) NMR spectra enabled the assignment of all protons for the hydrolysis products which supports structures I and II. The 1H NMR chemical shifts of the β -protons (4.83 and 4.46 ppm) of the serine residue of I were consistent with those found in pseudomycins A, B and C, indicating that the peptide macrocycle was intact. Furthermore, as expected, the TOCSY spectrum did not show the typical amide proton as part of the serine spin system. On the other hand, in II the serine β -protons underwent considerable upfield shifts (3.78 and 3.74 ppm) suggesting that these protons were not bearing the lactone functionality. This and the fact that the β protons, in addition to the α proton, correlated to an amide

proton at 8.04 ppm in the TOCSY spectrum indicated that the lactone of the macrocycle rearranged to a peptide core as depicted in II.

 $\begin{tabular}{llll} \bf Table & \bf I \\ & \begin{tabular}{llll} \bf I \\ & \begin{tabular}{lllll} \bf I \\ & \begin{tabular}{llll} \bf I \\ & \begin{tabular}{llll}$

Amino Acid	Position	I	II
Ser	NH	-	8.04
	α	4.30	4.30
	β1	4.83	3.78
	β2	4.46	3.74
Dab-1b	NH	9.19	7.99
	α	4.06	4.19
	β1	2.03	2.15
	β2		2.01
	γ1	3.03	2.92
	γ2	2.96	
Asp	NH	8.51	8.20
	α	4.61	4.56
	β1	2.89	2.84
	<u>β2</u>	2.83	2.75
Lys	NH	7.90	8.11
Bys	α	4.23	4.06
	<u>β1</u>	1.79	1.76
	β2	1.71	1.68
		1.27	1.30
	<u>γ1</u>		1.25
	γ2	1.54	1.54
	δ	2.84	2.84
	<u>8</u>	7.34	7.34
Dab-2 ^b	NH ₂ NH	8.35	8.31
Dab-2	α	4.29	4.34
		2.14	2.09
	β1	1.98	1.91
	β2	2.90	2.92
	γ	7.53	7.49
Thr	NH ₂ NH	7.73	7.74
TIII		4.24	4.21
	α	3.98	3.98
	β	1.18	1.16
	Υ	1 1.10	1.10

10

Table I (continued)

Amino Acid	Position	I	II
Dhb	NH	9.65	9.26
	β	6.69	6.62
	γ	1.69	1.66
OHAsp	NH	7.82	7.83
	α	4.95	4.99
	β	4.72	4.75
ClThr	NH	7.92	7.95
	α	4.90	4.62
	β	4.27	4.25
	γ1	3.48	3.57
	γ2	3.42	3.51

a Chemical shifts reported are relative to solvent signal (1.94 ppm).
b Assignments may be interchanged.

Other pseudomycin or syringomycin compounds having an N-acyl group may be deacylated using the same general procedures described above.

- 1. A process for deacylating an N-acyl side-chain of a pseudomycin comprising the step of reacting a pseudomycin with a deacylating enzyme selected from the group consisting of ECB deacylase and polymyxin acylase to produce a pseudomycin nucleus.
- 2. The process of Claim 1 wherein said pseudomycin nucleus is represented by either structure I or II

10

ΝH2 II

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

The process of Claim 1 wherein said pseudomycin is 3. selected from the group consisting of pseudomycin A, A', B, B', C, and C'.

15

5

10

4. A compound having the following structure

or a pharmaceutically acceptable salt, hydrate or solvate thereof, prepared by the process of Claims 1, 2 or 3.

5. A compound having the following structure

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

- 5 A pseudomycin nucleus prepared by reacting a pseudomycin with a deacylating enzyme selected from the group consisting of ECB deacylase and polymyxin acylase.
- 7. The pseudomycin nucleus of Claim 6 wherein said 10 pseudomycin is selected from the group consisting of pseudomycin A, A', B, B', C, and C'.
 - A process for deacylating an N-acyl side-chain of a syringomycin comprising the step of

reacting a syringomycin matural-product with a deacylating enzyme selected from the group consisting of ECB deacylase and polymyxin acylase to produce a syringomycin nucleus.

The process of Claim 7 wherein said syringomycin 9. nucleus is represented by either structure III or IV

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

- 10. A syringomycin nucleus prepared by reacting a syringomycin with a deacylating enzyme selected from the group consisting of ECB deacylase and polymyxin acylase.
- 10 11. A compound having the following structure

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

12. A compound having the following structure

$$H_2N$$
 H_2N
 H_2N

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

Please □	type a	plus	sign	(+) ins	side this	box	+	

Approved for use through 9/30/98, OMB 0651-0032

		Patent and Tr	ademark C	office: U.S. DEF	PARTMENT OF	COMMERCE
		Attorney Docket I	Number	X-1165	60	
DECLARATION FO	R [First Named Inve	ntor	Adam	Joseph Kreu	zman
UTILITY OR DESIG	N		COMPL	ETE IF KNO)WN	
PATENT APPLICATION	ON .	Application Numl	ber			
		Filing Date				
X Declaration Submitted with Initial Filing	,	Group Art Unit				
Declaration Submitted after Initial Filing		Examiner Name				
As a below named inventor, I hereby declare t	hat					
		low port to my pame				
My residence, post office address, and citizenshi believe I am the original, first and sole Inventor			rininal first	and joint invent	or (if plural name	es are listed
believe I am the original, lifst and sole inventor	I for which a paten	t is sought on the Inve	ntion entitle	ed:	(ii piarai nam	o uro iloto
PROCESS F	OR DEACYL	ATION OF LIPO	DEPSI	PEPTIDES		
the specification of which						
is attached hereto OR						
X was filed on 08 Ju (MM/DD/YYYY)	ine 2000 a	s United States Applica	ation Numb	er or PCT Inter	national	
Application PCT/US00/15018 Number	and was amend (MM/DD/YYYY		12	2 July 2001	(if applicab	le).
II hereby state that I have reviewed and understa	nd the contents of	f the above-identified s	pecification	, including the o	claims, as amen	ded by any
amendment specifically referred to above. I acknowledge the duty to disclose information w	high is material to	notontohility as definer	1 in Title 37	Code of Feder	al Regulations &	1.56
acknowledge the duty to disclose information w	mon is material to	paternasinty de definition				
I hereby claim foreign priority benefits under Title Inventor's certificate, or § 365(a) of any PCT inte America, listed below and have also identified be	rnational application	on which designated at the box, any foreign ap	least one of the least	country other th r patent or inver	an the United St	ates of
	untry	Foreign Filing I	Date	Priority Not Claimed	Certified Cop	y Attached NO
Number(s)		(WINDD/TTT	'	T		
					L.J.	
Additional foreign application numbers ar	e listed on a suppl	emental priority sheet	attached he	ereto:		
I hereby claim the benefit under Title 35, United	States Code § 119	e) of any United State	es provision	nal applications(s) listed below.	
Application Number(s) 60/143,968		Date (MM/DD/YYYY) 15 July 1999		numbers	I provisional app are listed on a si leet attached hei	upplemental

Please	type a	plus	sign	(+)	inside	this	box

Approved for use through 9/30/98. OMB 0651-0032 Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

			_				_	
n	_	\sim	1	AR	ΛΤ	П	$\boldsymbol{\cap}$	NI
	-	۱.		AK	-			IV

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s), or § 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

Mulcu pecame available perme	en the ming date of the prior application	Taric the heading of the international	
U.S. Parent	PCT Parent	Parent Filing Date	Parent Patent Number
Application Number	Number	(MM/DD/YYYY)	(if applicable)

Additional U.S. or PCT international application numbers are listed on a supplemental priority sheet attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent

and Trademark Office connected therewith:	
Attorney Name	Reg. No.
Arvie J. Anderson	45,2 <u>63</u>
Lynn D. Apelgren	45,341
Robert A. Armitage	27,417
Brian P. Barrett	39,597
Michael T. Bates	34.121
Roger S. Benjamin	27.025
Gary M. Birch	48,881
William R. Boudreaux	35,796
Steven P. Caltrider	36,467
Paul R. Cantrell	36,470
Charles E. Cohen	34,565
Donald L. Comeglio	30,741
Gregory A. Cox	47,504
Paula K. Davis	47,517
Elizabeth A. Dawalt	44,646
John C. Demeter	30,167
Manisha A. Desai	43,585
Joanne Longo Feeney	35,134
Paul J. Gaylo	- 36,808
Francis O. Ginah	44,712
Janet A. Gongola	48,436
Amy E. Hamilton	33,894
Frederick D. Hunter	26,915
Thomas E. Jackson	33,064
Charles Joyner	30,466
Gerald P. Keleher	43,707

Additional registered practitioner(s) named on a supplemental sheet attached hereto.

Attorney Name	Reg. No.
James J. Kelley	41,888
Paul J. Koivuniemi	31,533
Robert E. Lee	27,919
Kirby Lee	47,744
James P. Leeds	35,241
Nelsen L. Lentz	38,537
Douglas K. Norman	33,267
Arleen Palmberg	40,422
Thomas G. Plant	35,784
Edward Prein	37,212
Grant E. Reed	_41,264
James J. Sales	33,773
Michael J. Sayles	32,295
Robert L. Sharp	45,609
David M. Stemerick	40,187
Mark J. Stewart	43,936
Robert D. Titus	_40,206
Robert C. Tucker	45 <u>,165</u>
Tina M. Tucker	47,145
MaCharri Vorndran-Jones	36,711
Gilbert T. Voy	43,972
Thomas D. Webster	39,872
Lawrence T. Welch	29,487
Alexander Wilson	45,782
Dan L. Wood	48,613

Direct all correspondence to:											
Name	E	LI LILLY AND	COMPAN	<u> </u>							
Address	P	TTN: TINA M.	TUCKER								
Address	ī	ILLY CORPO	RATE CEA	ITER/DO	31104	-0					
City	Į į	NDIANAPOLIS		State		NDIANA		ZIP	4	6285	
Country			Те	lephone	•		(317) 27		Fax	(317)-276-	
I hereby declar believed to be punishable by	e true; a	all statements mad nd further that thes imprisonment, or l of the application	se statements ooth, under S	s were ma ection 100	de with 01 of Ti	the knowledq tie 18 of the U	ge that willful	false statemer	nts and tr	ne like so made	are
Name of Sole or First Inventor:							ed for this u	nsigned inv	entor	·	
Given	Adam				ddle	Joseph	Family	Kreuzn	nan	Suffix e.g. Jr.	
Name	Adan	1		Nai	me ,		Traine			1 0.3.0	
-			em JE	seph	7/	ugmat			Date	1 / /	oj .
Name	ignature		im Je		Kia	W LN	Country	USA	Date		O) USA
Name Inventor's S	ignature	Greenwood		State	Kia	1 100 10 10 11		USA	Date	11/29/	
Name Inventor's S Residence:	ignature City	Greenwood 4183 Willow	w Wind E	State	Kia	1 100 10 10 11		USA	Date	11/29/	
Name Inventor's Si Residence:	ignature City	Greenwood 4183 Willow	w Wind E	State	Kia	1 100 10 10 11		USA	Date	11/29/	
Inventor's Si Residence: Address Post Office	ignature City	Greenwood 4183 Willow SAME AS A	w Wind E	State	Kia	1 100 10 10 11		USA	Date	11/29/ Citizenship	



$\mathcal{O}_{\mathcal{A}}$	
F	

Please type a plus sign (+) inside this box	+

П		ليط			Е	Patent a	A and Tradem	pproved for us ark Office: U.	e through	gh 9/30/98. ARTMENT	OMB 06	551-0032 MMERCE
				ECL				<u> </u>				
Name of A	ddition	al Joint Inventor, if an	y:			AP	etition has	s been filed	for this	unsigne	ed inver	ntor
Given	i didinappan			Middle Name			Family Name	Kulanth	aivel		uffix .g. Jr.	
Name Inventor's Signature	Dai	omiappan 10		mlle	ela	Ø,			Date	11/2	-910	1
Residence: (City C	armel	Stat	e	NIA	\mathcal{I}^{T}	Country	USA		Citizensl	nip U	SA
Address		14907 Admiral Way			٠۴٠	`						
Post Office A	ddress	SAME AS ABOVE										
City Car	mel		Stat	e IN	IN zip 46032 Country USA						******	
Γ			T					· · · · ·				
Name of A	ddition	al Joint Inventor, if an	у:			AP	etition ha	s been filed	for this	s unsigne	ed inve	ntor
Given Name	Micha	el		Middle Name ~	Johi	n Family Name		Rodriguez		Suffix e.g. Jr.		
Inventor's Signature	ON.	had John John	7	-					Date	11/2	19/01	
Residence:	City In	dianapolis	<i>V</i>	State	IN	√ Co	untry US	SA		Citizens	hip U	SA
Post Office A	Address	7649 Gordonshire (Court		1	<u> </u>						
Post Office A	Address	SAME AS ABOVE				_						
city Ind	ianapo	olis	Stat	e IN	Zip	46	278	Country	US	Α		
Name of A	Addition	al Joint Inventor, if an	y:		A Peti	ition h	as been f	iled for this	unsign	ed inven	tor	
Given				Middle Name			Family Name				Suffix e.g. Jr.	
Inventor's Signature								C	ate			
Residence:	City			Stat	:e	Co	ountry			Citizens	hip	
Post Office												
Post Office		SAME AS ABOVE										_
City			State		Zip			Country				
			T									- :
Name of A	Addition	nal Joint Inventor, if an	ıy:		A Peti	ition h	as been f	iled for this	unsign	ned inven		
Given Name	-			Middle Name			Family Name				Suffix e.g. Jr.	
Inventor's Signature									Date			
Residence:	City			State		C	ountry			Citizens	ship	
Post Office												
Post Office		SAME AS ABOVE	,			Γ'		 	-1			
City			Stat	te	Zip			Country				